arXiv:1302.2428v1 [hep-ph] 11 Feb 2013

CAS-KITPC/ITP-349

Hydrodynamics and Transport Coefficients in an IR-deformed Soft-Wall AdS/QCD Model at Finite Temperature

SHINGO TAKEUCHI and YUE-LIANG WU

State Key Laboratory of Theoretical Physics
Kavli Institute for Theoretical Physics China (KITPC)
Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

Abstract

We extend an IR-deformed soft-wall AdS/QCD model at zero temperature to a model at finite temperature and perform hydrodynamics. To have the in-falling boundary condition to make the hydrodynamic analysis possible, we treat the infrared energy scale factor in our metric as a temperature-depending parameter. Then by carrying out the hydrodynamic analysis, we compute the transport coefficients, the diffusion constant and the shear viscosity, through the linear response theory.

1 Introduction

To carry out the analyses in the strongly coupled region would be regarded as the key of many important unsolved problems in the contemporary high energy theoretical physics. One of these problems will be the low energy dynamics of quantum chromodynamics(QCD). Actually, many ideas in the non-perturbative analysis of QCD have been motivated. Among these, there are three interesting classes, chiral dynamic model based on large N_c [1], the lattice gauge theory [2] and holography [3, 4] (gauge/gravity correspondence, AdS/QCD, etc).

In the lattice gauge theory, Minkowski space-time in the original theory is replaced with a finite volume Euclidian discretized space-time with analytic continuation of time direction to imaginary time direction. As a result, the degree of freedom of theories becomes finite, and the non-perturbative numerical analysis for the action itself, which is Monte Carlo simulation, becomes available. However, Monte Carlo simulation has a notorious problem named as the sign problem, when fermions are involved in the Monte Carlo. Currently, no mean for it has been invented yet, and the current analyses in the lattice are always carried out by getting around it [5].

On the other hand, it is well known that the holography is a duality between the strongly coupled field theories and the weakly coupled gravities. It actually originates in the superstring theory in which the quantized open and closed strings at low energy can be identified with particles known in field theories and gravities [6]. As a result, behaviors of the low energy open strings describe the supersymmetric gauge theory. Then as an important matter in the correspondence, $U(N)$ supersymmetric gauge theory arises on N overlapped D-branes from the open strings sticking to these N overlapped D-branes at low energy. On the other hand, D-branes can be identified with black branes in the supergravity [7].

Thus one has two ways to describe a low energy D-branes. As a result, it is known that one can conjecture the duality between p-dimensional large-N supersymmetric $SU(N)$ gauge theory with large 't Hooft coupling and near-horizon geometry of black $p+1$-branes in the condition that the quantum effect of gravity and the length of string can be neglected. Although there is no exact proof for this correspondence at this moment, it is particularly expected that the duality between the $\mathcal{N} = 4$ four-dimensional large-N $SU(N)$ supersymmetric gauge theory [3, 4] and the five-dimensional anti de Sitter space is valid.

One of the great advantages in the gauge/gravity correspondence compared with the lattice gauge theory would be that it is irrelevant to the problem arising when one involves fermions like the sign problem. Because the space-time in the gauge/gravity correspondence is not discretized space, and the main analysis are carried out in the weakly coupled gravity side.

However, the current gauge/gravity correspondence also has a problem. It is that the dual gauge theories are always non-realistic as long as the gravity side is a solution (Top-down model), and gauge theories become supersymmetric. Due to this, many results in the field theory side in the current gauge/gravity correspondence is the ones independent of detail of theories or no more than qualitative ones or results just in supersymmetric model. On the other hand, once getting away from the study based on a solution in the gravity side, constructing holographic models in bottom-up way is also conducted energetically [8–10].

Anyway, the point that the gauge/gravity correspondence can be irrelevant to the peculiar problem in the treatment of fermions in the lattice gauge theory would be one of the great advantages. For this reason, the low energy dynamics of QCD has recently been studied intensively in the framework of the gauge/gravity correspondence. This came to the motivation on an IR-deformed soft-wall AdS/QCD model [*] [11], which is a kind of the holographic bottom-up model [9].

Recently, we have made the extension of an IR-deformed AdS/QCD model [11] to the finite temperature system [12, 13], where the deformed bulk vacuum and potential term have to be introduced for the scalar field to satisfy the equations of motion. This is because if one straightforwardly extends the model to the finite temperature, the solution of the equation of motion diverges for the dilaton. Only by deforming the bulk vacuum and potential term, one can obtain the smooth solution for the dilation. With such a treatment, we have examined the critical temperature of chiral symmetry breaking [12] through the analysis of the quark number susceptibility, and the meson spectrum [13]. In the analysis of the meson spectrum, we have carried out the numerical analysis of equation of motion for the fluctuations on the bulk gravity. Such a numerical analysis for the mass spectrum has been considered as a basic method of analyses in holographic QCD as well as the hydrodynamics.

One of the crucial points of the hydrodynamics is that it can be considered to be independent of detail of theories, and so independent of the transport coefficients, where the transport coefficients are the ones defined in the framework of the hydrodynamic analysis. In particular, the ratio η/s between the shear viscosity (η) and the entropy density (s) is the quantity characterizing the actual QCD. For this reason, the ratio η/s has been examined very much in the gauge/gravity correspondence [14]. Besides, the holographic hydrodynamics has played an important role in the longstanding problem in the causal hydrodynamics [15]. and quark-gluon plasma described by Bjorken flow [16].

With such a circumstances, turning to the hydrodynamics, we are going to work out in this paper the holographic hydrodynamics in an IR-deformed AdS/QCD model at finite temperature studied in [12, 13]. Some interesting features mentioned above will be investigated in the future.

---

[*] We have referred to this IR-deformed soft-wall AdS/QCD model as a predicted model in our former papers [12, 13].

Then we would like to mention the organization of this paper. In Chap.2, we are going to review the hydrodynamics and the transport coefficients obtained from linear response theory. In Chap.3, we will introduce our holographic model, and show the way how the model is extended to the finite temperature system. It is shown that if one takes the same way as the one in ref. [13], it turns out that the analysis becomes too complicated to be carried out. Therefore, in this paper, we will propose an alternative way, which is simpler than the one in ref. [13] as the quark-gluon plasma is considered to be above the critical temperature. It will be seen that there are in general four options in the numerical calculations for some factors, while only one of them is physically acceptable. In Chap.4, we will sort out the notation used in this paper. In Chap.5, we will carry out the hydrodynamic analysis for the fluctuation of the balk gravity dual to $U(1)$ baryon current in the scalar mode. In the analysis, a necessity to cancel the divergence at the horizon arises as usual. As our model involves the dilaton which makes the analysis complicated. However, we will show how to contain it by exploiting the integral constant. By using GKP-W relation [4], we will read out the retarded Green function for $U(1)$ baryon current in the scalar mode and its diffusion constant. In Chap.6, we are going to carry out the hydrodynamic analysis for the fluctuation of the balk gravity dual to $SU(2)$ flavor current in the vector mode as well as in Chap.5. Again by using the GKP-W relation [4], we will read out the retarded Green function. In Chap.7, we shall carry out the hydrodynamic analysis for the fluctuation of the tensor mode in the balk gravity dual to the energy-momentum tensor. Thus by using the GKP-W relation [4], we can directly read out the viscosity with the retarded Green function and evaluate the ratio η/s. Our summary and conclusions will be presented in Chap.8.

2 Brief Review on Hydrodynamics and Linear Response Theory

To begin with, we would like to describe the basic matters in hydrodynamics and linear response theory used in this paper. The description in this chapter is basically following the review papers [17].

The hydrodynamics is an effective theory to describe the macroscopic dynamics at large distances and time-scales. Conserved quantities are considered to survive in such large distances and time-scales, and the energy-momentum tensor $T^{\mu\nu}$ is one of conserved quantities. The hydrodynamics is formulated by the hydrodynamic equation instead of action principle. The hydrodynamic equation for the energy-momentum tensor is given as

$$\partial_\mu T^{\mu\nu} = 0, \tag{2.1}$$

where

$$T^{\mu\nu} = (\varepsilon + P)u^\mu u^\nu + P g^{\mu\nu} + \tau^{\mu\nu} \tag{2.2}$$

with energy density ε, pressure P, local fluid velocity u^μ and the $\tau^{\mu\nu}$ given as

$$\tau^{\mu\nu} = -\eta\left(\partial^\mu u^\nu + \partial^\nu u^\mu - \frac{2}{3}\eta^{\mu\nu}\partial_\alpha u^\alpha\right) - \zeta\eta^{\mu\nu}\partial_\alpha u^\alpha. \tag{2.3}$$

Here η and ζ mean shear and bulk viscosities, respectively. In a curved space, it is given as

$$\tau^{\mu\nu} = -P^{\mu\alpha}P^{\nu\beta}\left\{\eta\left(\nabla_\alpha u_\beta + \nabla_\beta u_\alpha - \frac{2}{3}g_{\alpha\beta}\nabla u\right) - \zeta g_{\alpha\beta}\nabla u\right\}, \tag{2.4}$$

where $P^{\mu\nu} = g^{\mu\nu} + u^\mu u^\nu$. In the above, η and ζ are regarded as the transport coefficients. In what follows we consider the fluid in the rest frame, $u^\mu = (1, 0, 0, 0)$.

In eq.(2.2), substituting $u^\mu = (1, 0, 0, 0)$ and expanding as $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ ($\eta_{\mu\nu} = \mathrm{diag}(-1, 1, 1, 1)$ and $h_{\mu\nu}$ means fluctuations), it turns out that $\delta\langle\tau^{xy}\rangle$ is given as

$$\delta\langle T^{xy}\rangle = i\omega\eta h_{xy}, \tag{2.5}$$

where we have represented the formula in the momentum space. It is known from the linear response theory that the response of an operator $\mathcal{O}$ for the external field $\phi_{(0)}(k)$ is given as

$$\delta\langle\mathcal{O}(k)\rangle = -G_{\mathrm{R}}^{\mathcal{O}\mathcal{O}}\mathcal{O}(k), \tag{2.6}$$

where $G_{\mathrm{R}}^{\mathcal{O}\mathcal{O}}(k) \equiv -i\int_{-\infty}^{\infty} d^4x e^{-ikx}\langle[\mathcal{O}(t, \boldsymbol{x}), \mathcal{O}(0, 0)]\rangle\theta(t)$ is the retarded Green function. Then by comparing eq.(2.5) with eq.(2.6), we can obtain the Kubo formula with regard to the shear viscosity as

$$\eta = -\lim_{\omega\to 0}\frac{1}{\omega}\,\mathrm{Im}\,G_{\mathrm{R}}^{xy\ xy}, \tag{2.7}$$

where $G_{\mathrm{R}}^{xy\ xy}(k) \equiv -i\int_{-\infty}^{\infty} d^4x e^{-ikx}\langle[T^{xy}(t, \boldsymbol{x}), T^{xy}(0, 0)]\rangle\theta(t)$.

Next we turn to the diffusion in the conserved current and the energy-momentum tensor. Toward a conserved current j^μ with $0 = \partial_\mu j^\mu$, the constitutive equation for the conserved current is given as

$$\boldsymbol{j} = \rho\boldsymbol{u} - D\nabla j^t = -D\nabla j^t, \tag{2.8}$$

where $j^t = \rho$ and D mean the charge density and the diffusion constant respectively, and we have taken account of the fluid rest frame. Then the Fick's low is valid as

$$\partial_t\rho - D\nabla^2\rho = 0. \tag{2.9}$$

This gives the following dissipation relation for the charge density,

$$\omega = -iDk^2, \tag{2.10}$$

which will be the pole in the retarded Green function in the charge density, $G_{\mathrm{R}}^{tt}(k) \equiv -i\int_{-\infty}^{\infty} d^4x e^{-ikx}$ $\langle[\rho(t, \boldsymbol{x}), \rho(0, 0)]\rangle\theta(t)$. Next, we consider the diffusion in the the energy-momentum tensor T^{ti} with $i = x, y$.

First, from the constitutive equation in the curved space, one can obtain

$$T^{zi} = -\frac{\eta}{\varepsilon + P}\partial_z T^{ti} = -\eta \partial_z u^i. \tag{2.11}$$

Here, we have arranged the k along with the x^3-axis as $k = (0, 0, k)$ for simplicity. Then, from $\partial_\mu T^{\mu i} = 0$, one can obtain

$$\partial_t T^{ti} - \frac{\eta}{\varepsilon + P}\partial_z^2 T^{ti} = 0. \tag{2.12}$$

This gives the dissipation relation in the energy-momentum tensor T^{ti},

$$\omega = -i\frac{\eta}{\varepsilon + P}k^2, \tag{2.13}$$

which will be the pole in the retarded Green function $G_{\mathrm{R}}^{ti\ ti}$.

Finally, we write down the decomposition of the energy-momentum current and the $U(1)$ current under the little group $SO(2)$ toward i-direction ($i = x, y$):

scalar mode: T_{00}, T_{03}, T_{33}, ${T^i}_i$ and J_0, J_3,

vector mode: T_{0i}, T_{3i} and J_i,

tensor mode: $T_{ij} - \delta_{ij}{T^k}_k/2$.

3 An IR-deformed AdS/QCD Model at Finite Temperature

We will start with the following geometry, which is deformed from Schwarzschild AdS_5 black hole geometry in the IR-region by a factor μ_g as

$$ds^2 = a^2(z)\left(-f(z)dt^2 + \sum_{i=1}^{3} dx_i^2 + \frac{dz^2}{f(z)}\right) \tag{3.1}$$

with $a^2(z) = (1/z^2 + \mu_g^2)/l^2$ and $f(z) = 1 - (z/z_0)^4$ (z_0 means the location of the horizon) and we have put the AdS radius l as 1 in what follows. The coordinate z is in the relation with the usual coordinate r as $z = 1/r$. This geometry has Hawking temperature $T = 1/(\pi z_0)$, and is asymptotically AdS_5 space-time.

We will consider the following $U(1) \times SU_L(2) \times SU_R(2)$ soft-wall model with scalar field on the background (3.1) as

$$S = \int d^5x\,\sqrt{-g}e^{-\Phi(z)}\left(R + \frac{12}{l^2}\right) + \int d^5x\,\sqrt{-g}e^{-\Phi(z)}\Bigg[-\frac{1}{4g_{\mathrm{U(1)}}^2}F_{MN}F^{MN}$$
$$+\mathrm{Tr}\left\{-\frac{1}{4g_{\mathrm{SU(2)}}^2}\left(F_{L,MN}F_L^{MN} + F_{R,MN}F_R^{MN}\right) + |D_M X(z)|^2 - m_X^2|X(z)|^2 - \frac{\lambda}{4}|X(z)|^4\right\}\Bigg], \tag{3.2}$$

where $m_X^2 = -3$ [8,9] and $l = 1$. The trace is performed for the $SU(2)$ algebra mentioned in what follows. We write F_{MN}, $F_{L,MN}$ and $F_{R,MN}$ as $F_{MN} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu$ and $F_{L,MN} \equiv \partial_\mu B_{L,\nu} - \partial_\nu B_{L,\mu} - i[B_{L,\nu}, B_{L,\mu}]$, where $B_{L,\mu} = B^a_{L,\mu} t^a$ with the $SU(2)$ Lie algebra t^a $(a = 1, 2, 3)$, and now we have skipped to describe for R part. As for the dual operators for these in the boundary theory, for example see the table in ref. [8]. Using these, the covariant derivative can be written as $D_M X = \partial_M X + i(B_{L,M} X - X B_{R,M})$.

The bulk vacuum of the scalar field is given as

$$X(z) = \frac{v(z)}{2}\, \mathbf{1}_2, \tag{3.3}$$

where $\mathbf{1}_2$ means 2×2 unit matrix, and $v(z)$ is given in Table.1. As it is known that $v(z)$ behaves around the boundary as

$$v(z) = m_q \zeta\, z + \frac{\sigma}{\zeta} z^3 + \mathcal{O}(z^5). \tag{3.4}$$

From AdS/CFT correspondence, m_q and σ can be interpreted as the quark mass and quark condensate, respectively. As for ζ, see Table.1 and 2.

Model	$v(z)$	Parameters
IIb	$z(A + Bz^2)(1 + Cz^4)^{-5/8}$	$A \equiv m_q\zeta$, $B \equiv \sigma/\zeta$, $C \equiv (B^2/(\mu_d\gamma^2))^{4/5}$

Table 1: These are taken from ref. [11]. The numerical values for these are given in Table.2, which are fixed by minimizing the breaking of the Gell-Mann-Oakes-Renner relation, $f_\pi^2\ m_\pi^2 = 2m_q\ \sigma$, at the 1% level and the experimental values: $m_\pi = 139.6$ MeV and $f_\pi = 92.4$ MeV.

Model	λ	m_q (MeV)	$\sigma^{\frac{1}{3}}$ (MeV)	γ	μ_g	μ
IIb	0	4.07	272	0.112	257	0.205
IIb	9	6.79	229	0.20	257	0.205

Table 2: The numerical values for Table.1. These are taken from [11]. $\mu \equiv \mu_g/(2\pi T)$ with $T = 0.2$GeV, $\zeta = \sqrt{3}/(2\pi)$ and $\mu_g = \sqrt{3}\mu_d$. Here we should notice that we will treat μ_g as a temperature-depending parameter which is different from the consideration in our previous paper [13] and will be discussed below in this chapter. The numerical values in this Table are taken from our previous paper [13].

Here we would like to touch on the field theory dual to our model (3.2) with the geometry (3.1). Although it is not an exact statement as our model is a bottom-up model and does not stand on the configuration of D-branes, the assumed dual field theory would be $D = 1 + 3$ SU(N_c) gauge theory with $U(1)$ baryon symmetry and $SU(N_f)_L \times SU(N_f)_R$ chiral flavor symmetry as global symmetries at large 't Hooft coupling at finite temperature, where $N_f = 2$ and the large-N_c limit is taken, and no charge is concerned for the global symmetries in the model.

We have shown in [12] that the quark number susceptibility blows up when temperature is around $160 \sim 190$ MeV, which is related to the chiral symmetry breaking/restoration. We have further examined the mass spectra for the vector and axial-vector mesons as the function of temperature in ref. [13], which has suggested that the critical temperature locates at around 200 MeV.

The temperature and the entropy density in the dual field theory are given from Hawking temperature and Bekenstein-Hawking formula as

$$T_H = \frac{1}{\pi z_0} \quad \text{and} \quad s = \frac{a^3(z_0)}{4G_5} = \frac{N_c^2}{2\pi}\left\{(\pi T)^2 + \mu_g^2\right\}^{3/2}, \tag{3.5}$$

where we have used the relation $G_5 = \pi/(2N_c^2)$. The relation of $g_{U(1)}$ and $g_{SU(2)}$ to the soft-wall model are given as [9]

$$g_{U(1)}^2 = \frac{16\pi^2}{N_c^2} \quad \text{and} \quad g_{SU(2)}^2 = \frac{12\pi^2}{N_c}. \tag{3.6}$$

The equation for the dilaton can be obtained from the equation of motion for the scalar field as

$$\Phi'(z) = \frac{3a'(z)}{a(z)} + \frac{\left(f(z)v'(z)\right)'}{f(z)v'(z)} - \frac{a^2(z)}{f(z)v'(z)}\left(m_X^2 v(z) + \frac{\lambda}{2}v^3(z)\right). \tag{3.7}$$

It will be noticed that in our previous paper [13], we have fixed $X(z)$ with considering a regularized term $v_1(z)\ln f(z)$ as $X(z) = \frac{1}{2}\Big\{v_1(z)\ln f(z) + v_0(z)\Big\}\,\mathbf{1}_2$ (and in accordance with this extra term, the coupling parameter λ has to be modified), where $v_0(z)$ in this equation corresponds to $v(z)$ in this paper. The explicit form for $v_1(z)$ is referred to ref. [13]. As it can be seen that without considering the extra term $v_1(z)\ln f(z)$, the dilaton $\Phi'(u)$ in the vicinity of the horizon starts with the order $(u-1)^{-1}$ as

$$\Phi'(u) = \frac{\Phi_{-1}(T,\mu)}{u-1} + \mathcal{O}(1) \tag{3.8}$$

with the numerators,

$$\begin{aligned}\Phi_{-1}(T,\mu) = &-\frac{1}{A\left(6C(\mu)z_0^4-4\right)-2Bz_0^2\left(C(\mu)z_0^4+6\right)}\Big[A\Big\{m_X^2\left(4\mu^2+1\right)\left(C(\mu)z_0^4+1\right)-6C(\mu)z_0^4+4\Big\}\\ &+Bz_0^2\Big\{m_X^2\left(4\mu^2+1\right)\left(C(\mu)z_0^4+1\right)+2\left(C(\mu)z_0^4+6\right)\Big\}\Big],\end{aligned} \tag{3.9}$$

where $u \equiv z^2/z_0^2$ and $\mu \equiv \mu_g/(2\pi T)$ (Later they are defined at eq.(4.11) and (4.12)), and A, B and $C(\mu)$ are given in Table.1. Then it turns out that the contribution $\Phi_{-1}(T,\mu)/(u-1)$ appears at the order $(u-1)^{-1}$ in the equation of motion for the fluctuations at the vicinity of the horizon. Therefore, the contribution $\Phi(\mu)$ appears at the order $(u-1)^{-2}$ in the equation of motion. This prevents the solutions of the fluctuations from taking the in-falling boundary condition, where the in-falling boundary condition is determined at the order $(1-u)^{-2}$ in the equations of motion. For this reason, it has been found in [13] to be useful for introducing the extra term $v_1(z)\ln f(z)$ in order to regularize the divergence and obtain the in-falling boundary condition healthily.

However, for our present purpose in this paper, it turns out that the extra term $v_1(z)\ln f(z)$ makes the hydrodynamic analysis very complicated. Thus we have to discard the technique adopted in [13] with introducing the extra term $v_1(z)\ln f(z)$ and find an alternative way. A simple idea is to consider the energy scale μ (or μ_g) as a temperature-depending parameter $\mu(T)$, so that one can adjust the parameter μ to make the numerator $\Phi_{-1}(T,\mu)$ vanishes at the order $(1-u)^{-1}$. As a result, we obtain the μ which vanishes the numerator $\Phi_{-1}(T,\mu)$ at each temperatures as shown in Fig.1, and finally we can take the in-falling boundary condition.



Figure 1: We plot the positive and real values of μ found as the function of temperature. They are obtained from vanishing the numerator $\Phi_{-1}(T,\mu)$ at the order $(u-1)^{-1}$ in eq.(3.9). As a consequence, one can healthily take the in-falling boundary condition. The left and right plots are different just in the scale of x-axis. The lower curve (red points) and upper curve (blue points) in each figure represent two branches of solution for a given temperature.

In Fig.1, it is noticed that there are two branches for the solution, which are plotted in each figure by the lower curve (red) and upper curve (blue) for a given temperature. Then the question is which branch is physically meaningful. Before answering the question, we may make a comment on the relation of μ in our previous paper [13] as shown in the rightmost column of table.2 and in this paper given in Fig.1.

For the result of the model IIb, two branches start at $T_c = 0.2$ GeV, and the value of μ at the point for the branch of the lower curve (red points) matches also mostly the same value of μ shown in the rightmost column of table.2. Then, let us consider which branches we should take.

First, we shall recall that μ is the factor appearing in the factor $a^2(z)$ in eq.(3.1) as $a^2(z) = 1/z^2 + \mu_g^2 = 1/z^2 + (2\pi T\mu)^2$. Then, one can see that if μ remains finite at high temperature, as temperature increases, our bulk space-time becomes completely different from the Schwarzschild AdS_5 black hole space-time. It means that if μ does not vanish as temperature goes up, the gauge/gravity correspondence becomes invalid. On the other hand, if μ vanishes at high temperature, as temperature increases, our bulk space-time goes back to the Schwarzschild AdS_5 blackhole space-time. At that time, since our model (3.2) has the extra terms (dilaton, scalar field and gauge fields) other than the Einstein-Hilbert action, the dual field theory is not the $\mathcal{N} = 4$ supersymmetric gauge theory even if the background geometry is the Schwarzschild AdS_5 blackhole space-time. However, what the background geometry goes back to the Schwarzschild AdS_5 blackhole space-time in the high temperature limit would be a consistent condition as a holographic AdS/QCD model. In this sense, eventually, the physically acceptable branch would be the one represented by the lower curve (red points), and the rest would be inconsistent.

An interesting point is that the effect of μ does not disappear abruptly, but gradually disappears as temperature increases. The effect of μ is a specific one in our AdS/QCD model, and it would be interesting to study its phenomenon in the future. Further, it would also be interesting that the branch of the model IIb starts at about $T = 0.2$ GeV, where the temperature $T_c = 0.2$ GeV is roughly the critical temperature for the Hadron/Plasma transition.

4 Preliminaries for Hydrodynamic Analysis

Before making a hydrodynamic analysis, let us first clarify the indices used in this paper

$$M, N = x_0,\ x_1,\ x_2,\ x_3,\ z \text{ or } 0,\ 1,\ 2,\ 3,\ 4 \tag{4.1}$$

$$\mu, \nu = x_0,\ x_1,\ x_2,\ x_3 \text{ or } 0,\ 1,\ 2,\ 3 \tag{4.2}$$

$$i, j, k = x_1,\ x_2 \text{ or } 1,\ 2 \tag{4.3}$$

$$\alpha, \beta = x_1,\ x_2,\ x_3 \text{ or } 1,\ 2,\ 3 \tag{4.4}$$

$$a, b, c = 1\ ,2\ ,3 \text{ (indices for } SU(2) \text{ algebra)} \tag{4.5}$$

where x_0 is coordinate of time, x_1, x_2 and x_3 are coordinates of space on the boundary and z is radial direction of the bulk.

We are going to examine the retarded Green function for the scalar mode of $U(1)$ baryon current, the vector mode of $SU(2)$ flavor current and the shear viscosity from the tensor mode of the energy-momentum tensor through AdS/CFT correspondence. For this purpose, we will analyze the hydrodynamics in the bulk gravity of the fluctuations,

$$g_{MN} \longrightarrow G_{MN} = g_{MN} + h_{MN}, \tag{4.6}$$

$$g^{MN} \longrightarrow G^{MN} = g^{MN} - h^{MN}, \tag{4.7}$$

$$A_M \longrightarrow B_{0M} = 0 + A_M, \tag{4.8}$$

$$B^a_M \longrightarrow B^a_{0M} = 0 + B^a_M, \tag{4.9}$$

where the background of the gauge fields are vanishing. Without confusing, we simply use the same character for the total gauge fields and the fluctuation of gauge fields. These fluctuations are in linear order in equations of motion. In this paper, we will treat the dilaton as the static background.

We show the classification of fluctuations in the $SO(2)$ little group for x and y-directions as

$$\begin{aligned}
&\text{Scalar mode: } h_{00},\, h_{03},\, h_{33},\, h_{zz},\, h_{0z},\, h_{3z},\, h^k_{\ k} \quad \text{and} \quad A_0,\, A_3,\, A_z, \\
&\text{Vector mode: } h_{0i},\, h_{3i},\, h_{zi} \quad \text{and} \quad A_i, \\
&\text{Tensor mode: } h_{ij} - \delta_{ij} h^k_{\ k}/2.
\end{aligned} \tag{4.10}$$

One may see the correspondence of the above classification with the one given in Sec.2.

In AdS/CFT correspondence, the fluctuations of the bulk gauge field A_μ acts as the source for the R-charge current J_μ on the dual field theory. Since it is known that R-charge can be regarded as an analog of the $U(1)$ baryon number in AdS/CFT correspondence, we will consider that A_μ acts as the source for the global $U(1)$ baryon charge current J_μ (For example, see ref. [21]). Further, $B_{L,\mu}$ and $B_{R,\mu}$ also act as the source for the global $SU(N_f)_L \times SU(N_f)_R$ chiral flavor currents where $N_f = 2$ in this paper. The gravitational perturbations in the bulk act as the source for the stress-energy tensor $T_{\mu\nu}$ on the dual theory.

In our following analysis, for convenience we will define the normalized radial coordinate

$$u \equiv \frac{z^2}{z_0^2}, \tag{4.11}$$

and the normalized frequency, momentum and the factor

$$\mathfrak{w} \equiv \frac{\omega}{2\pi T}, \quad \mathfrak{q} \equiv \frac{q}{2\pi T} \quad \text{and} \quad \mathfrak{u} \equiv \frac{\mu_g}{2\pi T}. \tag{4.12}$$

Finally, let us mention the gauge fixing condition in our present consideration. Like other papers (For example, ref. [22]), we choose the axial gauge condition and the Landau gauge as follows

$$A_z = 0, \quad B^a_z = 0 \quad \text{and} \quad h_{Mz} = 0, \tag{4.13}$$

$$\partial^\mu A_\mu = 0 \quad \text{and} \quad \partial^\mu B_{L,\mu} = \partial^\mu B_{R,\mu} = 0, \tag{4.14}$$

where the Landau gauge is imposed in the four-dimensional space-time on the boundary.

5 Analysis on the $U(1)$ baryon current

We shall first perform hydrodynamic analysis in the scalar mode shown in Table.(4.10) of $U(1)$ baryon current. To begin with, we can write the equation of motion for the fluctuation of A_M in terms of the field strength as

$$\partial_M \left(\sqrt{-g(u)} e^{-\Phi(u)} F^{MN}(u) \right) = 0, \tag{5.1}$$

where $g(u) \equiv \det g_{MN}(u)$. We will carry out analysis in the momentum space by performing the plane wave expansion as

$$A_\mu(u, x_0, x_3) = \int \frac{dw\, dq}{(2\pi)^2} e^{-i\omega x_0 + iqx_3} A_\mu(u, \omega, q)\,, \tag{5.2}$$

where we have rotated the direction of momentum to x_3, and pay attention to the same notation used in the gauge fields before and after Fourier transformation.

$$\omega g^{00}(u) A_0'(u) - q g^{33}(u) A_3'(u) = 0\ , \tag{5.3}$$

$$\partial_u \left(e^{-\Phi(u)} \sqrt{-g(u)} g^{00}(u) g^{uu}(u) A_0'(u) \right) = e^{-\Phi(u)} \sqrt{-g(u)} g^{00}(u) g^{33}(u) \left(\omega q A_3(u) + q^2 A_0(u) \right), \tag{5.4}$$

$$\partial_u \left(e^{-\Phi(u)} \sqrt{-g(u)} g^{33}(u) g^{uu}(u) A_3'(u) \right) = e^{-\Phi(u)} \sqrt{-g(u)} g^{00}(u) g^{33}(u) \left(\omega q A_0(u) + w^2 A_3(u) \right), \tag{5.5}$$

$$\partial_u \left(e^{-\Phi(u)} \sqrt{-g(u)} g^{\alpha\alpha}(u) g^{uu}(u) A_\alpha'(u) \right) = e^{-\Phi(u)} \sqrt{-g(u)} g^{\alpha\alpha}(u) \left(g^{00} \omega^2 A_\alpha(u) + g^{33} q^2 A_\alpha(u) \right), \tag{5.6}$$

where $' \equiv \partial_u$.

From eqs.(5.3) and (5.4), the equation of motion for A_t' can be obtained as follows

$$\left(\frac{z_0}{2}\right)^2 \frac{d}{du} \left[\frac{\partial_u (e^{-\Phi(u)} \sqrt{-g(u)} g^{00}(u) g^{uu}(u) A_0'(u))}{e^{-\Phi(u)} \sqrt{-g(u)} g^{00}(u) g^{33}(u)} \right] - \left(\frac{g^{00}(u)}{g^{33}(u)} \mathfrak{w}^2 + \mathfrak{q}^2 \right) A_0'(u) = 0. \tag{5.7}$$

For a technical reason, we will perform a rescaling as

$$-\frac{2}{z_0^2} e^{-\Phi(u)} \sqrt{-g(u)} g^{00}(u) g^{uu}(u) A_0'(u) \equiv \mathcal{A}_0'(u), \tag{5.8}$$

where we attach the factor $-2/z_0^2$ in the front to make the leading term of the rescaling as unit in the expansion around $u = 0$. Note that when taking $\mu = 0$, it recovers the case of Schwarzschild AdS_5 black hole geometry. With the above rescaling, the equation of motion is given by

$$\left(\frac{z_0}{2}\right)^2 \frac{d}{du} \left[\frac{\partial_u \mathcal{A}_0'(u)}{e^{-\Phi(u)} \sqrt{-g(u)} g^{00}(u) g^{33}(u)} \right] = \frac{\frac{g^{00}(u)}{g^{33}(u)} \mathfrak{w}^2 + \mathfrak{q}^2}{e^{-\Phi(u)} \sqrt{-g(u)} g^{00}(u) g^{uu}(u)} \mathcal{A}_0'(u). \tag{5.9}$$

To make the hydrodynamic analysis, let us formally write down the solution to the first order hydrodynamics as follows

$$A_0'(u) = C_0^{(0)}(u) \frac{e^{\Phi(u)}}{\sqrt{1 + 4\mu^2 u}} (1 - u)^{-i\mathfrak{w}/2} \left(1 + \mathfrak{w} F_1^{(0)}(u) + \mathfrak{q}^2 G_1^{(0)}(u) + \cdots \right), \tag{5.10}$$

where $C_0^{(0)}(u)$, $F_1^{(0)}(u)$ and $G_1^{(0)}(u)$ are going to be fixed below. It is noticed that the in-falling boundary condition has been used to yield the factor $(1 - u)^{-i\mathfrak{w}/2}$ in the solution. This is realized by taking the special solution of μ, as shown in Fig.1, with requiring the numerator in the contribution at the order of

$(u-1)^{-1}$ to be vanishing in the dilaton $\Phi'(u)$. Then we can see readily that $C_0^{(0)}$ can be treated as a constant,

$$C_0^{(0)}(u) \equiv C_0^{(0)}, \tag{5.11}$$

and the rest coefficients are obtained by the following equations,

$$\frac{z_0^2}{8}\partial_u \left\{ \frac{iC_0^{(0)} + 2(1-u)\partial_u F_1^{(0)}(u)}{(1-u)e^{-\Phi(u)}\sqrt{-g(u)}g^{00}(u)g^{33}(u)} \right\} = 0, \tag{5.12}$$

$$\frac{z_0^2}{4}\partial_u \left(\frac{\partial_u G_1^{(0)}(u)}{e^{-\Phi(u)}\sqrt{-g(u)}g^{00}(u)g^{33}(u)} \right) = \frac{C_0^{(0)}}{e^{-\Phi(u)}\sqrt{-g(u)}g^{uu}(u)g^{00}(u)}, \tag{5.13}$$

where the former and later can be obtained from the order at $\mathfrak{w}$ and $\mathfrak{q}^2$ in eq.(5.10). Let us solve the above equations by taking the following three steps:

1. performing the integration on the whole equations to get rid of the derivative $\partial_u\{\cdots\}$ or $\partial_u(\cdots)$,
2. rewriting the equations in the form "$\partial_u F_1^{(0)} = \cdots$" or "$\partial_u G_1^{(0)} = \cdots$",
3. carrying out the integration over the whole equations to obtain $F_1^{(0)}$ and $G_1^{(0)}$,

we then obtain the coefficients as follows

$$F_1^{(0)}(u) = C_0^{(0)}\left(F_B^{(0)} - \frac{2F_H^{(0)}e^{-\Phi(0)}\log(u)}{z_0^2} \right) - C_0^{(0)}\left(\frac{i}{2} - \frac{2F_H^{(0)}e^{-\Phi(0)}\left(\Phi'(u)|_{u=0} - 2\mu^2\right)}{z_0^2} \right) u + \mathcal{O}(u^2), \tag{5.14}$$

$$G_1^{(0)}(u) = C_0^{(0)}\left(G_B^{(0)} - \frac{2F_H^{(0)}e^{-\Phi(0)}\log(u)}{z_0^2} \right) + C_0^{(0)}\left(1 + \frac{2G_H^{(0)}e^{-\Phi(0)}\left(\Phi'(u)|_{u=0} - 2\mu^2\right)}{z_0^2} \right) u + \mathcal{O}(u^2) \tag{5.15}$$

with

$$C_0^{(0)} = -\frac{\mathfrak{q}z_0^2(A_0^{(0)}\mathfrak{q} + A_3^{(0)}\mathfrak{w})}{2(G_H\mathfrak{q}^2 + F_H\mathfrak{w})}. \tag{5.16}$$

Here $C_0^{(0)}$ has been fixed by using eq.(5.4) with the Dirichlet boundary condition that the boundary value of A_0 and A_3 are given by $A_0^{(0)}$ and $A_3^{(0)}$. Note that if the full integration in step 1 or 3 is difficult, instead of it, one can first carry out the Tyler expansion around $u=0$, and after that perform integration toward its low order terms. As we will use eventually the GKP-W relation, the resulting solutions are the ones around the boundary. Where $F_H^{(0)}$, $F_B^{(0)}$, $G_H^{(0)}$ and $G_B^{(0)}$ are the integral constants. $F_H^{(0)}$ and $G_H^{(0)}$ appear from the integration in step 3, and they can be taken arbitrarily. On the other hand, $F_B^{(0)}$ and $G_B^{(0)}$ appear from the integration in step 1, and they are in general fixed such that the solutions do not diverge at $u=1$. Where the subscripts "H" and "B" denote the relevant constants which are fixed from the horizon and the boundary, respectively.

To put it more concretely, after step 2, one can see that $F_1^{(0)\prime}(u)$ and $G_1^{(0)\prime}(u)$ behave as $F_1^{(0)\prime}(u)\,, G_1^{(0)\prime}(u) \sim 1/(u-1)$ around $u=1$, which generally leads to the solution with logarithmic divergence, such a solution is ill-defined. To obtain a physically meaningful solution, a simple way is to vanish these contributions by exploiting the integral constants $F_H^{(0)}$ and $G_H^{(0)}$, so that the numerators become zero.

In the actual calculation, $F_H^{(0)}$ can easily be fixed to be

$$F_H^{(0)} = -\frac{iC_0^{(t)} e^{\Phi(1)} z_0^2}{2\sqrt{1+\mu^2}}. \tag{5.17}$$

while fixing G_H is in general complicated as the integrating becomes difficult due to the dilaton. For this reason, we have to perform integration around $u=1$ to fix the integral constant G_H. It will be shown that as long as choosing the integral constant G_H appropriately, we can arrive at the needed solution which has no logarithmic divergence and becomes well-defined in the whole bulk. Let us conduct step 1 with expansion around $u=1$

$$\begin{aligned}
&\frac{\partial_u G_1^{(0)}(u)}{e^{-\Phi(u)}\sqrt{-g(u)}g^{00}(u)g^{33}(u)} \\
&= \int du \frac{C_0^{(0)}}{e^{-\Phi(u)}\sqrt{-g(u)}g^{uu}(u)g^{00}(u)} \\
&= \int du \Big\{ C_0 + C_1(u-1) + C_2(u-1)^2 + C_3(u-1)^3 + \cdots \Big\} \\
&= G_H^{(0)} + C_0 u + C_1\left(\frac{u^2}{2} - u\right) + C_2\left(\frac{u^3}{3} - u^2 + u\right) + C_3\left(\frac{u^4}{4} - u^3 + \frac{3}{2}u^2 - u\right) + \cdots \\
&= G_H^{(0)} + \sum_{n=0}^{\infty}\frac{(-1)^n}{n+1}C_n + C_0 t + C_1\left(\frac{t^2}{2} - t\right) + C_2\left(\frac{t^3}{3} - t^2 + t\right) + C_3\left(\frac{t^4}{4} - t^3 + \frac{t^2}{2} - t\right) + \cdots
\end{aligned} \tag{5.18}$$

with

$$C_n \equiv \frac{C_0^{(0)}}{n!}\,\partial_u^{(n)}\left(\frac{1}{e^{-\Phi(u)}\sqrt{-g(u)}g^{uu}(u)g^{00}(u)}\right)\Bigg|_{u=1} \quad \text{and} \quad t \equiv u-1, \tag{5.19}$$

where "(n)" means the number of derivative with regard to u. In the above equations, we have performed an expansion from the second to the third line and written its result in symbolic way. From the third to the forth line, we have explicitly written the integral constant $G_H^{(0)}$. From the forth to the fifth line, we have changed the variable u to $t \equiv u-1$. It is seen that when we rewrite it in the form that "$\partial_u G_1^{(0)} = \cdots$", there is no term with $1/(u-1)$ in the expansion beyond the constant term. Therefore, we can readily obtain the expression of $G_H^{(0)}$ to be

$$G_H^{(0)} = -\sum_{n=0}^{\infty}\frac{(-1)^n}{n+1}C_n \equiv \frac{C_0^{(0)} z_0^2}{2e^{-\Phi(1)}\sqrt{1+4\mu^2}} G_A\big(\mu, \{\Phi^{(n)}\}\big), \tag{5.20}$$

where we have rewritten symbolically the concise expression for convenience in the following actual calculation. Here $G_A\left(\mu, \{\Phi^{(n)}\}\right)$ is a function with $\left\{\Phi^{(n)}\right\} \equiv \left\{\Phi^{(1)}(u)|_{u=1}, \Phi^{(2)}(u)|_{u=1}, \Phi^{(3)}(u)|_{u=1}, \cdots\right\}$, when $\mu = 0$ and $\{\Phi^{(n)}\} = \{0\}$, one has $G_A(0, \{0\}) = 1$. In this paper, $G_H^{(0)}$ will be treated symbolically.

It can be shown that the solutions near the boundary behave as follows

$$A_0'(u) = -\frac{2C_0^{(0)}}{z_0^2}\left(G_B^{(0)}\mathfrak{q}^2 + F_B^{(0)}\mathfrak{w}\right)\log(u) + C^{(0)}e^{\Phi(0)}\left(1 + G_B^{(0)}\mathfrak{q}^2 + F_B^{(0)}\mathfrak{w}\right), \tag{5.21}$$

$$A_3'(u) = \frac{2C_0^{(0)}}{z_0^2}\mathfrak{w}\left(G_B^{(0)}\mathfrak{q} + F_B^{(0)}\frac{\mathfrak{w}}{\mathfrak{q}}\right)\log(u) - C_0^{(0)}e^{-\Phi(0)}\frac{\mathfrak{w}}{\mathfrak{q}}\left(1 + G_B^{(0)}\mathfrak{q}^2 + F_H^{(0)}\mathfrak{w}\right). \tag{5.22}$$

The boundary action S_0 is given from the on-shell action in the quadratic order of fluctuation as

$$S_0 = \frac{1}{g^2 z_0^2}\int \frac{dw\,dq}{(2\pi)^2}\left\{\left(1 + \frac{4\mu^2\sqrt{u}}{z_0}\right)A_0(u)A_0'(u) - \left(1 + \frac{4\mu^2\sqrt{u}}{z_0}\right)\sum_\alpha A_\alpha(u)A_\alpha'(u)\right\}. \tag{5.23}$$

By adopting the prescription for obtaining the retarded Green function in AdS/CFT correspondence [18], we arrive at the following two-point retarded green function of $U(1)$ baryon charge current in the scalar mode

$$G_{(\mathrm{R})}^{00} = \frac{N^2T^2}{8}\frac{e^{\Phi(0)-\Phi(1)}\sqrt{4\mu^2+1}k^2}{2i\pi T\omega - G_0(\mu, \{\Phi^{(n)}\})k^2}, \tag{5.24}$$

$$G_{(\mathrm{R})}^{33} = \frac{N^2T^2}{8}\frac{e^{\Phi(0)-\Phi(1)}\sqrt{4\mu^2+1}\omega^2}{2i\pi T\omega - G_0(\mu, \{\Phi^{(n)}\})k^2}. \tag{5.25}$$

From these results, we can read out the diffusion constant (2.9) as

$$D = \frac{G_A(\mu, \{\Phi^{(n)}\})}{2\pi T}. \tag{5.26}$$

6 Analysis on $SU(2)$ vector and axial-vector flavor currents

Let us make hydrodynamic analysis for the vector mode shown in Table (4.10) of $SU(2)$ vector current and axial-vector flavor current. To begin with, let us combine the $SU(2)$ gauge fields into the vector field V_M and the axial-vector field W_M

$$V_M(u) \equiv \frac{1}{2}(B_{L,M}(u) + B_{R,M}(u)) \quad \text{and} \quad W_M(u) \equiv \frac{1}{2}(B_{L,M}(u) - B_{R,M}(u)). \tag{6.1}$$

The equations of motion for the vector field V_x^a and the axial-vector field W_x^a are given as

$$\mathrm{V}\ :\ D_{V,M}\left(\sqrt{-g(u)}e^{-\Phi(u)}\,F_V^{a,MN}(u)\right) = 0, \tag{6.2}$$

$$\mathrm{AV}\ :\ D_{W,M}\left(\sqrt{-g(u)}e^{-\Phi(u)}\,F_W^{a,MN}(u)\right) + \ g_{SU(2)}^2 e^{-\Phi(u)}\sqrt{-g(u)}\ v^2(u)g^{MN}(u)W_M^a(u) = 0, \tag{6.3}$$

where $v(u)$ has been defined in eq.(3.3), and "V" and "AV" mean the field strength consisting of the vector and the axial-vector fields, respectively. $F_{V,MN}^a(u)$ and $D_{V,M}(u)$ are dictated as $F_{V,MN}^a(u) \equiv \partial_M V_N^a(u) - \partial_N V_M^a(u) + if^{abc}V_M^b(u)V_N^c(u)$ (f^{abc} is $SU(2)$ structure constant), and $D_{V,M} \equiv \partial_M(u) - iV_M(u)$ ($F_{W,MN}(u)$ and $D_{W,M}(u)$ are similar). It is seen that eq.(6.2) can be obtained from eq.(6.3) by dropping the term proportional to $g^2_{SU(2)}$. Thus our main task here will focus on solving eq.(6.3).

Now let us write down the equation of motion (6.3) to the linear order, and its result is given by

$$\partial_u\big(e^{-\Phi(u)}\sqrt{-g(u)}g^{\alpha\alpha}(u)g^{uu}(u)W^{a\prime}_\alpha(u)\big) = e^{-\Phi(u)}g^{\alpha\alpha}(u)\big(g^{00}(u)\omega^2 + g^{33}(u)k^2 - g^2_{SU(2)}v^2(u)\big)W^a_\alpha(u). \tag{6.4}$$

which cannot simply be solved by the double integral dictated above eq.(5.14) due to the extra term proportional to $g^2_{SU(2)}v^2(u)$ in the potential part. Thus we may factorize it as follows

$$W^a_\alpha(u) \to \rho(u)\mathcal{W}^a_\alpha(u). \tag{6.5}$$

Then eq.(6.4) becomes the following form

$$\partial_u\Big(e^{-\Phi(u)}\sqrt{-g(u)}g^{\alpha\alpha}(u)g^{uu}(u)\mathcal{W}^{a\prime}_\alpha(u)\Big) + \rho(u)\left\{\Omega(u) - e^{-\Phi(u)}\sqrt{-g}\rho g^{\alpha\alpha}(u)(g^{00}\omega^2 + g^{33}k^2)\right\} = 0 \tag{6.6}$$

with

$$\Omega(u) \equiv \partial_u\Big(e^{-\Phi(u)}\sqrt{-g(u)}g^{\alpha\alpha}(u)g^{uu}(u)\rho'(u)\Big) - g^2_{SU(2)}e^{-\Phi(u)}\sqrt{-g(u)}\rho(u)g^{\alpha\alpha}(u)v^2(u). \tag{6.7}$$

which indicates that when $\rho(u)$ satisfies $\Omega(u) = 0$, it becomes available to obtain the solution $W^a_\alpha(u)$ from the double integral. Nevertheless, to fully solve the equation $\Omega(u) = 0$ is technically difficult. Practically, considering that the needed solution at the last stage is the one only in the vicinity of the horizon and the boundary, we shall obtain the solutions in the expansion around $u = 0$ and $u = 1$.

First, let us obtain the solution in the expansion around $u = 0$. For this purpose, expanding $\Omega(u)$ given in eq.(6.7), we arrive at the following form

$$\big(\alpha_0 + \mathcal{O}(u)\big)\rho''(u) + \big(\beta_0 + \mathcal{O}(u)\big)\rho'(u) + \Big(\frac{\gamma_0}{u} + \gamma_1 + \mathcal{O}(u)\Big)\rho(u) = 0 \tag{6.8}$$

with

$$\alpha_0 \equiv \frac{2}{z_0^2}, \quad \beta_0 \equiv -\frac{2}{z_0^2}\left(\Phi'(u)\big|_{u=0} - 2\mu^2\right), \quad \gamma_0 \equiv \frac{1}{2}g^2_{SU(2)}A^2, \quad \gamma_1 \equiv g^2_{SU(2)}A^2\left(Bz_0^2 + 3A\mu^2\right). \tag{6.9}$$

As a result, we obtain the general solution $\rho(u)$ as a linear combination of two confluent hypergeometric functions ${}_1F_1$ and U with two integral constants. We set two integral constants in such a way that ${}_1F_1$ vanishes and the leading of $\rho(u)$ in the expansion of $u = 0$ becomes 1. Eventually, we obtain $\rho(u)$ satisfying eq.(6.9) to the expanded order as

$$\rho(u) = \mathrm{Exp}\left(-\frac{\Delta + \beta_0}{2\alpha_0}u\right)\Gamma\left(1 - \frac{\gamma_0}{\Delta}\right)U\left(-\frac{\gamma_0}{\Delta}, 0, \frac{\Delta}{\alpha_0}u\right) \tag{6.10}$$

with $\Delta \equiv \sqrt{(\beta_0)^2 - 4\alpha_0\gamma_1}$. Where the functions $\Gamma(u)$ and $U(u)$ denote the gamma function and the confluent hypergeometric function, respectively.

Now we shall try to solve $\Omega(u) = 0$ in the vicinity of $u = 1$. It turns out that a general solution is given by a linear coupling of Confluent Hypergeometric Function U and Laguerre Polynomials L_n. However, it finds difficult to fix the integral constants analytically, such that the constants in the solution at $u = 1$ becomes common with the ones at $u = 0$ of eq.(6.10) due to the lack of information for the solution of $\rho(u)$ in the interior domain of the bulk. Therefore, we will only treat the solution $\rho(u)$ around $u = 1$ symbolically.

Having obtained $\rho(u)$ satisfying $\Omega(u) = 0$ as in eq.(6.10), we can solve the equation of motion by using the double integral similar to the previous chapter. With the results of μ given in Table 1, we can formally write down the solution of $\mathcal{W}^a_\alpha$ in the hydrodynamic expansion with regard to $\mathfrak{w}$ and $\mathfrak{q}^2$ as

$$\mathcal{W}^a_\alpha = C^{(a)}_\alpha(u)(1-u)^{-i\mathfrak{w}/2}\left(1 + \mathfrak{w}F^{(a)}_1(u) + \mathfrak{q}^2 G^{(a)}_1(u) + \cdots\right). \tag{6.11}$$

With the same way stated above eq.(5.14), the coefficients can be written as

$$C^{(a)}_\alpha(u) = \frac{\mathcal{W}^{a(0)}_\alpha}{1 + G^{(a)}_B q^2 + F^{(a)}_B \omega}, \tag{6.12}$$

and

$$F^{(a)}_1(u) = F^{(a)}_B + \left(\frac{2F^{(a)}_H e^{\Phi(0)}}{\rho(0)^2} - \frac{iC^{(a)}_0}{2}\right)u + \mathcal{O}(u^2), \tag{6.13}$$

$$G^{(a)}_1(u) = G^{(a)}_B + \left\{C^{(a)}_0(\log(u) - 1) + \frac{2G^{(a)}_H e^{\Phi(0)}}{\rho^2(0)}\right\}u + \mathcal{O}(u^2), \tag{6.14}$$

where $\mathcal{W}^{a(0)}_\alpha$ denotes the boundary value of $\mathcal{W}^a_\alpha$, and $F^{(a)}_B$, $G^{(a)}_B$ and $F^{(a)}_H$, $G^{(a)}_H$ mean integral constants which are fixed at the boundary and the horizon as the same as the integral constants appeared in eq.(5.14) and (5.15). To be explicit, they are fixed as

$$F^{(a)}_H = \frac{1}{2}iC^{(a)}_0\sqrt{4\mu^2+1}\,\rho^2(1)e^{-\Phi(1)}, \tag{6.15}$$

$$G^{(a)}_H = -\frac{1}{2}C^{(a)}_0\sqrt{4\mu^2+1}\,e^{-\Phi(1)}G_W(\mu, \{\Phi^{(n)}\}), \tag{6.16}$$

where $G_W(\mu, \{\Phi^{(n)}\})$ is a function as the same as eq.(5.20) and $G_W(0, \{0\}) = 1$. From the above analysis, we can write down the solutions in the vicinity of $u = 0$ with taking the factor ρ

$$\begin{aligned} W^a_\alpha = \mathcal{W}^{a(0)}_\alpha\rho(0) + \mathcal{W}^{a(0)}_\alpha \Bigg\{ & \frac{4\rho(0)e^{\Phi(0)}\left(F^{(a)}_H\omega + G^{(a)}_H q^2\right) + \rho^2(0)\left(2q^2\log(u) - 2q^2 - i\omega\right)}{2\rho(0)\left(F^{(a)}_B\omega + G^{(a)}_B q^2 + 1\right)} \\ & + \left(\rho(1) + \frac{i\rho(0)}{2}\omega\right)\Bigg\} u + \mathcal{O}(u^2). \end{aligned} \tag{6.17}$$

It is noticed that V_α^a can simply be read from the solution of W_α^a by taking $g^2_{SU(2)} = 0$.

Finally, we come to consider the two-point retarded Green function in the dual field theory through GKP-W relation. For this purpose, the boundary action in quadratic order at $u = 0$ is needed. One can see that it is given as

$$S_0 = -\frac{2}{g^2_{SU(2)} z_0^2}\left(1 + \frac{4\mu^2\sqrt{u}}{z_0}\right)\int \frac{dw\, dq}{(2\pi)^2}\sum_\alpha W_\alpha^a(u) W_\alpha^{a\prime}(u). \tag{6.18}$$

which enables us to obtain the two-point retarded Green function in vector mode as

$$G^{\alpha\alpha}_{\mathrm{AV,R}} = \frac{iN^2T^2}{8}\rho(0)\left\{\frac{4\rho(0)e^{\Phi(0)}\left(F_H^{(a)}\omega + G_H^{(a)}q^2\right) + \rho(0)^2\left(2q^2\log(u) - 2q^2 - i\omega\right)}{2\rho(0)\left(F_B^{(a)}\omega + G_B^{(a)}q^2 + 1\right)} + \left(\rho(1) + \frac{i\rho(0)\omega}{2}\right)\right\}. \tag{6.19}$$

The two-point retarded Green function $G^{\alpha\alpha}_{\mathrm{V,R}}(u)$ for $V_\alpha^a(u)$ can simply be resulted from the $G^{\alpha\alpha}_{\mathrm{AV,R}}(u)$ given in above with vanishing $g^2_{SU(2)}$.

7 Analysis on the tensor mode

We are now going to carry out the analysis for the tensor mode of the fluctuation shown in Table (4.10) in the dual gravity, which allows us to calculate the shear viscosity (2.7) and its ratio to the entropy density in the dual field theory. For this purpose, we will start with the equation of motion for the bulk gravity, which can be generally given as

$$R_{MN} - \frac{1}{2}G_{MN}R + \Lambda G_{MN} = \kappa^2 T_{MN}, \tag{7.1}$$

where $\kappa^2 = 8\pi G_5$ (G_5 is given below eq.(3.5)), $\Lambda = -6/l^2$ (l denotes the curvature radius of the AdS space and is taken to be 1) and T_{MN} means the energy-momentum tensor. From eq.(3.1), it is given as

$$T_{MN} \equiv g_{MN}\mathcal{L} + g^{PQ}\left\{\frac{1}{g^2_{U(1)}}F_{PM}F_{QN} + \frac{1}{g^2_{SU(2)}}\mathrm{Tr}\Big(F_{L,PM}F_{L,QN} + F_{R,PM}F_{R,QN}\Big)\right\} - 2\mathrm{Tr}D_MXD_NX, \tag{7.2}$$

where $\mathcal{L}$ means the Lagranigian (3.2) except for the Einstein-Hilbert part. It is noticed that when substituting the background (3.1) into the above equation of motion for gravity, there is a deviation proportional to μ_g, which indicates that the background (3.1) is not an exact solution of the equation of motion for gravity as the back-reacted geometry is not considered here. Nevertheless, the effect from the back-reacted geometry was found to be small [20], and also it turns out that the deviated part appears only in the diagonal part of the equation of motion, while in our present treatment it involves only the off-diagonal part of the equation of motion, (x, y)-component, so the deviated part will be discarded in this paper.

To evaluate the viscosity, we will consider (x,y)-component in eq.(7.1). Then the energy-momentum tensor at the linear order becomes as $T_{xy} = \mathrm{Tr}\left(g^{uu}|\partial_u X|^2 - m_X^2|X|^2 - \frac{\lambda}{4}|X|^4\right) h_{xy}$, and the equation of motion for h_{xy} can be obtained as

$$0 = h''_{xy}(u) + \frac{g'_1(u)}{g_1(u)} h'_{xy}(u) + g_2(u) h_{xy}(u) \tag{7.3}$$

with

$$\begin{aligned}
g_1(u) &\equiv \frac{\sqrt{u} f(u)}{a(u)}, \\
g_2(u) &\equiv g_3(u) + \frac{z_0^2}{4uf(u)^2}\left(\omega^2 - k^2 f(u)\right), \\
g_3(u) &\equiv -\frac{1}{2ua(u)f(u)}\Big[16uf(u)a''(u) + 8a'(u)\big(2uf'(u) + f(u)\big) + a(u)\big(s2uf''(u) + f'(u)\big) \\
&\qquad -6z_0^2 a(u)^3 + \frac{\kappa^2 z_0^2 a^2(u)}{4uf(u)} T_{xy}(u)\Big].
\end{aligned} \tag{7.4}$$

Here $g_3(u)$ represents the constant part toward ω and k in the potential term $g_2(u)$. It is seen that $g_3(u)$ is the obstacle to utilize the double integral dictated above eq.(5.14), which is similar to the case of eq.(6.4). Thus we may adopt the same analysis as the one for eq.(6.4), namely exploiting the factorization,

$$h_{xy}(u) = \chi(u)\phi(u), \tag{7.5}$$

and vanishing the extra term $g_3(u)$.

Under the factorization (7.5), we can rewrite eq.(7.3) as

$$\big(g_1(u)\chi^2(u)\phi'(u)\big)' + g_1(u)\chi^2(u)\left\{\Psi(u) + \frac{z_0^2}{4uf(u)^2}\left(\omega^2 - k^2 f(u)\right)\right\}\phi(u) = 0 \tag{7.6}$$

with

$$\Psi(u) \equiv \chi''(u) + \frac{g'_1(u)}{g_1(u)}\chi'(u) + g_3(u)\chi(u). \tag{7.7}$$

Then $\chi(u)$ is obtained by requiring $\Psi(u)$ to satisfy $\Psi(u) = 0$. However, it can be shown that there is difficult to solve the equation $\Psi(u) = 0$ fully. Similarly to the previous chapter, as the needed solution eventually is the one in the vicinity of the horizon and the boundary, we may consider the solution of $\chi(u)$ only in the vicinities at $u = 0$ and $u = 1$. Nevertheless, from the reason mentioned below, one cannot obtain analytically the solution in the expansion around $u = 1$.

Then we will try to obtain the solution of $\chi(u)$ in the vicinity of $u = 0$. To this purpose, expanding the constant part $\Psi(u)$ around $u = 0$, we will consider the following equation

$$\begin{aligned}
\chi''(u) + \left(\frac{1}{u} - 2\mu^2\right)\chi'(u) + \Big[&-\frac{1}{u^2} + \frac{1}{2u}\left(40\mu^2 + \kappa^2 T'_{xy}(u)\big|_{u=0}\right) \\
&+\frac{1}{4}\Big\{-8 + \kappa^2\left(8\mu^2 T'_{xy}(u)\big|_{u=0} + T''_{xy}(u)\big|_{u=0}\right)\Big\}\Big]\chi(u) = 0.
\end{aligned} \tag{7.8}$$

To obtain the solution $\chi(u)$ in the vicinity of $u=0$, we may assume the following form as the solution of $\chi(u)$,

$$\chi(u) = \frac{\chi_{-1}}{u} + \chi_0 + \chi_1 u + \chi_L \, u \log(u) \quad \text{with} \quad \chi_{-1} \equiv 1/z_0^2, \tag{7.9}$$

where χ_0, χ_1 and χ_L are the coefficients to be determined further. While we have fixed $\chi_{-1} = 1/z_0^2$. It can be seen that if we assume the form of solution as in eq.(7.9), then χ_{-1} can in general be arbitrary (and each coefficients except for χ_{-1} can be fixed so as to satisfy eq.(7.8)). Thus one should fix χ_{-1} at the first place as shown in ref. [19]. By doing so, one can see that $\chi(u)$ plays the same role as the background metric $g_{\alpha\alpha}$ at the vicinity of $u=0$, where $g_{\alpha\alpha}(u)$ in the vicinity of $u=0$ behaves as $g_{\alpha\alpha}(u) = \frac{1}{z_0^2}\frac{1}{u} + \frac{4\mu^2}{z_0^2} + \mathcal{O}(u)$. As a consequence, we can put the source of $\phi(u)$ as $h^{(0)}{}^{y}_{x}(u)$ with $h^{(0)}{}^{y}_{x}(u)$ the boundary value.

By substituting $\chi(u)$ with the form in eq.(7.9) and solving the equations in each order of it, we can fix the coefficients as

$$\begin{aligned}
\chi_0 &= \frac{\chi_{-1}}{2}\left(44\mu^2 + \kappa_2 T'_{xy}(u)\big|_{u=0}\right), &(7.10)\\
\chi_1 &= \frac{\chi_{-1}}{24\left(36\mu^2 + \kappa_2 T'_{xy}(u)\big|_{u=0}\right)}\Bigg[3\log(u)\left(36\mu^2 + \kappa_2 T'_{xy}(u)\big|_{u=0}\right)\Big\{1728\mu^4 + 92\kappa_2\mu^2 T'_{xy}(u)\big|_{u=0}\\
&\quad +\kappa_2\left(T''_{xy}(u)\big|_{u=0} + \kappa_2\left(T'_{xy}(u)\big|_{u=0}\right)^2\right)\Big\} - 2\kappa_2\left\{6T'_{xy}(u)\big|_{u=0}\left(268\mu^4 + 5\kappa_2\mu^2 T'_{xy}(u)\big|_{u=0} - 2\right)\right.\\
&\quad \left. +3T''_{xy}(u)\big|_{u=0}\left(54\mu^2 + \kappa_2 T'_{xy}(u)\big|_{u=0}\right) + 2T^{(3)}_{xy}(0)\right\} - 16128\mu^6\Bigg], &(7.11)\\
\chi_L &= -\frac{\chi_{-1}}{8}\left\{1728\mu^4 + 92\kappa_2\mu^2 T'_{xy}(0) + \kappa_2\left(T''_{xy}(u))\big|_{u=0} + \kappa_2\left(T'_{xy}(u)\big|_{u=0}\right)^2\right)\right\}, &(7.12)
\end{aligned}$$

where we have obtained the solution of $\chi(u)$ to the order of u, which will become necessary in the analysis in what follows.

Now we come to discuss the solution $\chi(u)$ in the vicinity of $u=1$. From the expansion of the coefficients of eq.(7.8) around $u=1$, one can see that a general solution in the vicinity of $u=1$ is given as a linear coupling of Confluent Hypergeometric Function U and Laguerre Polynomials L_n. However, similar to the case of $\rho(u)$ in the previous chapter, it is difficult to fix the integral constants analytically, so that the constants are common between the ones at $u=1$ and $u=0$ (eq.(7.9) with the coefficients (7.10)), due to lack of the information of the solution of $\chi(u)$ in the interior domain of the bulk. Thus $\chi(u)$ around $u=1$ will be treated symbolically in the present consideration.

In the same way with the previous sections, we can obtain the hydrodynamic solution for the off-diagonal component of the fluctuation $h_{xy}(u)$ in eq.(7.6) with $\chi(u)$ determined above. Consequently, it can be written as

$$h_{xy}(u) = \chi(u)\phi(u) = C_0^{(\phi)}\,\chi(u)\,(1-u)^{-i\mathfrak{w}/2}\left(1 + \mathfrak{w}F_1^{(\phi)}(u) + \mathfrak{q}^2 G_1^{(\phi)}(u) + \cdots\right) \tag{7.13}$$

with

$$C_0^{(\phi)} = \frac{h^{(0)}{}^x_{\ y}}{1 + G_B^{(\phi)}\mathfrak{w} + F_B^{(\phi)}\mathfrak{q}^2}, \tag{7.14}$$

$$F_1^{(\phi)}(u) = F_B^{(\phi)} - \frac{i}{2}u + \frac{i}{2}\left(-\frac{1}{2} + \frac{F_H^{(\phi)}}{z_0^4\chi_{-1}}\right)u^2, \tag{7.15}$$

$$G_1^{(\phi)}(u) = G_B^{(\phi)} - u + \left\{\frac{2\chi_{-1}z_0\log(u)\left(\chi_0 - \mu^2\chi_{-1}\right) + \chi_{-1}z_0\left(\chi_0 - \mu^2\chi_{-1}\right)}{2(\chi_{-1})^2 z_0} + \frac{G_B^{(\phi)}}{2(\chi_{-1})^2 z_0^4}\right\}u^2, \tag{7.16}$$

Where $h^{(0)}{}^x_{\ y}$ has been mentioned above eq.(7.10). $F_H^{(\phi)}$, $F_B^{(\phi)}$ and $G_H^{(\phi)}$, $G_B^{(\phi)}$ represent the integral constants fixed at boundary and horizon, which are similar to the integral constants appeared in eq.(5.14) and (5.15). More explicitly, they are fixed as

$$F_H^{(\phi)} = \frac{1}{\sqrt{4\mu^2+1}}, \tag{7.17}$$

$$G_H^{(\phi)} = -\frac{1}{\sqrt{4\mu^2+1}}G_0^{(\phi)}\left(\mu, \{\Phi^{(n)}\}\right). \tag{7.18}$$

Here we have used $\chi_{-1} = 1/z_0^2$. It enables us to write down the solution of $h_{xy}(u)$ in the vicinity of $u = 0$ as

$$\begin{aligned} h_{xy}(u) = {} & \frac{h^{(0)}{}^x_{\ y}\chi_{-1}}{u} + h^{(0)}{}^x_{\ y}(\chi_0 - \mathfrak{q}^2\chi_{-1}) + \frac{h^{(0)}{}^x_{\ y}}{2z_0^4\chi_{-1}^2}\Big[iF_H^{(\phi)}\omega + q^2\Big\{G_H^{(\phi)} + \chi_{-1}z_0^4\left(\chi_0 - \mu^2\chi_{-1}\right)\Big\} \\ & + 2z_0^4 q^2\chi_{-1}\left(\chi_0 - \mu^2\chi_{-1}\right)\log(u)\Big]u + \mathcal{O}(u^2) \end{aligned} \tag{7.19}$$

which shows that the expression diverges at $u = 0$. However, such a divergence arises from the contribution of $\chi(u)$ given in eq.(7.10), and the solution of $\phi(u)$ itself has been obtained healthily as it can be seen from eq.(7.13). Such a situation is the same as the one discussed in ref. [19] and also other papers based on ref. [19].

Now that we have obtained the solution, let us evaluate the viscosity through Kubo formula. The boundary action S_0 is given by

$$S_0 = S_{\text{on-shell}} + S_{\text{GH}} + S_{\text{ct}}, \tag{7.20}$$

with

$$S_{\text{GH}} = \frac{1}{8\pi G_5}\int d^4x\sqrt{-g^{(4)}}K, \tag{7.21}$$

$$S_{\text{ct}} = -\frac{1}{8\pi G_5}\int d^4x\sqrt{-g^{(4)}}\left(\frac{3}{l} + \frac{l}{4}R^{(4)}\right), \tag{7.22}$$

where S_{GH} and S_{ct} denote the Gibbons-Hawking term and the counter term respectively, and $g^{(4)}_{\mu\nu}$ is the four-dimensional induced metric, and K and $R^{(4)}$ mean the extrinsic curvature and the curvature on the boundary toward our geometry (3.1), respectively. They are found to be

$$S_{\text{on-shell}} = \frac{1}{2\kappa^2 z_0} \int \frac{dw\, dq}{(2\pi)^2} e^{-\Phi(u)} \frac{\sqrt{u}}{a(u)^2} \Big\{ 4a'(u) f(u) h_{xy}(u)^2 - 3a(u) f(u) h_{xy}(u) {h_{xy}}'(u) \Big\}\Big|_{u=0}, \tag{7.23}$$

$$S_{\text{GH}} = \frac{1}{2\kappa^2 z_0} \lim_{u\to 0} \int \frac{dw\, dq}{(2\pi)^2} e^{-\Phi(u)} \frac{\sqrt{u}}{a(u)} \Big\{ h_{xy}(u)^2 f'(u) + 4 f(u) h_{xy}(u) {h_{xy}}'(u) \Big\}, \tag{7.24}$$

$$S_{\text{ct}} = \frac{1}{8\kappa^2} \lim_{u\to 0} \int \frac{dw\, dq}{(2\pi)^2} e^{-\Phi(u)} \frac{12 a(u)^2 f(u) + k^2 f(u) - \omega^2}{a(u)^2 \sqrt{f(u)}} h_{xy}^2(u). \tag{7.25}$$

The retarded Green function is obtained by using the GKP-W relation

$$G^{(\text{R})}_{xy\ xy}(\omega, k) = -\frac{N^2 T^2}{16} \Big\{ G_H^{(\phi)} k^2 + 2\pi T \left(i F_H^{(\phi)} \omega - \chi_{-1}^2 \right) \Big\}. \tag{7.26}$$

which allows us to evaluate the shear viscosity via Kubo formula

$$\eta = -\lim_{\omega\to 0} \frac{\text{Im}\left(G^{(\text{R})}_{xy\ xy}(\omega, k=0) \right)}{\omega} = \frac{\pi}{8} F_H N^2 T^3. \tag{7.27}$$

with this result, it is then not difficult to obtain the ratio

$$\frac{\eta}{s} = \frac{1}{4\pi \sqrt{1+4\mu^2} (1 + 16\pi^2 \mu^2 T^2)^{3/2}}, \tag{7.28}$$

where the entropy density is given in eq.(3.5).

The ratio η/s as the function of temperature is shown in Fig.2, where the left plots and right plots is just in the scale of x-axis. The two curves (upper red points and lower blue points) represent the results obtained by using the μ in Fig.1, and the each plots in Fig.2 and Fig.1 correspond each other. The dashed horizontal line represents the KSS bound, $1/(4\pi)$ [23]. It can be seen that there are two branches with regard to the solutions of μ. As it has been discussed in the Chap.3, the physically acceptable branch is given by the lower curves (red points) in the plots.

8 Conclusions and Remarks

In this paper, we have extended an IR-deformed AdS/QCD model presented in ref. [11] to finite temperature system by another way different from the one proposed in ref. [13], worked out the hydrodynamics and computed transport coefficients. In this extension, we have found several branches, and we have chosen the branch in which our model can be consistent as a holographic model.

The specific things in our model are the parameters μ_g, $g_{SU(2)}$, λ, the scalar field X and dilaton Φ. Among these, the ones entering to our analysis in this paper has been μ_g thorough $\mu \equiv \mu_g/(2\pi T)$ and dilaton. We have treated the effect of these in symbolic way. The rest things will enter if one considers the background dual to the system at the finite chemical potential. However, the analysis with the finite chemical potential might be too complected to perform the hydrodynamic analysis.



Figure 2: We plot η/s given in eq.(7.28) using μ given in Fig.1. The lefts and right plots are different just in the scale of x-axis, The red and blue points represent the results obtained by using the μ in Fig.1. These plots and the plots in Fig.1 correspond each other. The dashed horizontal line represents the KSS bound, $1/(4\pi)$ [23]. Among these plots, the result represented by the red points corresponds to the results concerning the μ we have chosen as physically acceptable one.

A further direction of this study will be the holographic description of Bjorken flow [24]. Bjorken Flow is an effective model having been invented to describe evolution of quark-gluon plasma produced in high energy collision experiment. This has the property of the ideal fluid, and diffuses with the form of distribution in the boost invariant way from a point in Minkowski space-time where a collision happens. it is considered that the quark-gluon plasma being produced and diffusing in RHIC or LHC, etc can be described by using Bjorken Flow. However, it is known that understanding of Bjorken Flow in the frame work of the field theory is hard due to its strong coupling and real-time evolution. In such a situation, the gauge/gravity correspondence would be an effective measure. For the relevant studies, we would like to refer reader to the reference in the review papers [16]. In the analysis of these, we can see that hydrodynamic analysis is necessity. Therefore, using the hydrodynamic analysis formulated in this paper, we are going to try to the Bjorken flow in our model.

Acknowledgments The author S.T. would like to thank Y. Matsuo, S. Okazawa, T. Azuma, Y. Imamura and J. P. Shock for useful discussions. This work is supported in part the National Nature Science Foundation of China (NSFC) under Grants No. 10975170, No. 10821504 and No. 10905084; and the Project of Knowledge Innovation Program (PKIP) of the Chinese Academy of Science.

References

[1] Y. Nambu, Phys. Rev. Lett. **4** (1960) 380.

G. 't Hooft, Nucl. Phys. B **72**, 461 (1974).

E. Witten, Annals of Physics, **128**, 363 (1980).

Y. B. Dai and Y. L. Wu, Eur. Phys. J. C **39** (2004) S1 [arXiv:hep-ph/0304075].

[2] K. G. Wilson, Phys. Rev. D **10**, 2445 (1974).

F. J. Wegner, J. Math. Phys. **12**, 2259 (1971).

M. Creutz, Acta Phys. Slov. **61**, 1 (2011) [arXiv:1103.3304 [hep-lat]].

[3] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200].

E. Witten, Adv. Theor. Math. Phys. **2**, 505-532 (1998). [hep-th/9803131].

[4] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109].

E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].

[5] For a recent review, see e.g., S. Ejiri, Prog. Theor. Phys. Suppl. **186**, 510 (2010) [arXiv:1009.1186 [hep-lat]], and references therein.

[6] A. Neveu and J. Scherk, Nucl. Phys. B **36**, 155 (1972).

T. Yoneya, Prog. Theor. Phys. **51**, 1907 (1974).

J. Scherk and J. H. Schwarz, Nucl. Phys. B **81**, 118 (1974).

[7] G. T. Horowitz and A. Strominger, Nucl. Phys. B **360**, 197 (1991).

[8] J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, Phys. Rev. Lett. **95**, 261602 (2005) [arXiv:hep-ph/0501128].

[9] A. Karch, E. Katz, D. T. Son and M. A. Stephanov, Phys. Rev. D **74**, 015005 (2006) [arXiv:hep-ph/0602229].

[10] For a recent review, see e.g., Y. Kim, I. J. Shin and T. Tsukioka, arXiv:1205.4852 [hep-ph].

Y. Kim and D. Yi, Adv. High Energy Phys. **2011**, 259025 (2011) [arXiv:1107.0155 [hep-ph]].

[11] Y. Q. Sui, Y. L. Wu, Z. F. Xie and Y. B. Yang, "Prediction for the Mass Spectra of Resonance Mesons in the Soft-Wall AdS/QCD with a Modified 5D Metric," Phys. Rev. D **81**, 014024 (2010) [arXiv:0909.3887 [hep-ph]].

Y. Q. Sui, Y. L. Wu and Y. B. Yang, Phys. Rev. D**83**, 065030 (2011).

[12] L. -X. Cui, S. Takeuchi and Y. -L. Wu, Phys. Rev. D **84**, 076004 (2011) [arXiv:1107.2738 [hep-ph]].

[13] L. -X. Cui, S. Takeuchi and Y. -L. Wu, JHEP **1204**, 144 (2012) [arXiv:1112.5923 [hep-ph]].

[14] For a review, see e.g., S. Cremonini, Mod. Phys. Lett. B **25**, 1867 (2011) [arXiv:1108.0677 [hep-th]].

[15] M. Natsuume and T. Okamura, Phys. Rev. D **77**, 066014 (2008) [Erratum-ibid. D **78**, 089902 (2008)] [arXiv:0712.2916 [hep-th]].

[16] M. Rangamani, Class. Quant. Grav. **26**, 224003 (2009) [arXiv:0905.4352 [hep-th]].

M. P. Heller, R. A. Janik and R. Peschanski, Acta Phys. Polon. B **39**, 3183 (2008) [arXiv:0811.3113 [hep-th]].

A. Bernamonti and R. Peschanski, Nucl. Phys. Proc. Suppl. **216**, 94 (2011) [arXiv:1102.0725 [hep-th]].

[17] P. Kovtun, D. T. Son and A. O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005) [hep-th/0405231].

M. Natsuume, "AdS/CFT duality learned by linear response theory" Unpublished article written in Japanese.

[18] D. T. Son and A. O. Starinets, JHEP **0209**, 042 (2002) [arXiv:hep-th/0205051].

[19] G. Policastro, D. T. Son and A. O. Starinets, JHEP **0209**, 043 (2002) [hep-th/0205052].

[20] J. P. Shock, F. Wu, Y. L. Wu and Z. F. Xie, JHEP **0703** (2007) 064 [arXiv:hep-ph/0611227].

Y. L. Wu and Z. F. Xie, JHEP **0710** (2007) 009 [arXiv:0705.2360 [hep-ph]].

[21] S. Nakamura, Y. Seo, S. -J. Sin and K. P. Yogendran, Prog. Theor. Phys. **120**, 51 (2008) [arXiv:0708.2818 [hep-th]].

S. Kobayashi, D. Mateos, S. Matsuura, R. C. Myers and R. M. Thomson, JHEP **0702**, 016 (2007) [hep-th/0611099].

[22] M. Fujita, K. Fukushima, T. Misumi, M. Murata, Phys. Rev. **D80**, 035001 (2009). [arXiv:0903.2316 [hep-ph]].

M. Fujita, T. Kikuchi, K. Fukushima, T. Misumi, M. Murata, Phys. Rev. **D81**, 065024 (2010). [arXiv:0911.2298 [hep-ph]].

[23] P. Kovtun, D. T. Son and A. O. Starinets, JHEP **0310**, 064 (2003) [hep-th/0309213].

P. Kovtun, D. T. Son and A. O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005) [hep-th/0405231].

[24] J. D. Bjorken, Phys. Rev. D **27**, 140 (1983).